

# *Drosophila* divalent metal ion transporter *Malvolio* is required in dopaminergic neurons for feeding decisions

E. Søvik<sup>†,‡</sup>, A. LaMora<sup>†,1</sup>, G. Seehra<sup>†</sup>,  
A. B. Barron<sup>§</sup>, J. G. Duncan<sup>¶</sup>  
and Y. Ben-Shahar<sup>†,\*</sup> 

<sup>†</sup>Department of Biology, Washington University, St. Louis, MO, USA, <sup>‡</sup>Department of Science and Mathematics, Volda University College, Volda, Norway, <sup>§</sup>Department of Biological Sciences, Macquarie University, Sydney, NSW, Australia, and <sup>¶</sup>Department of Pediatrics, Washington University School of Medicine, St. Louis, MO, USA

<sup>1</sup>Present address: Department of Biology, McKendree University, Lebanon, IL 62254, USA

\*Corresponding author: Y. Ben-Shahar, Department of Biology, Washington University, 1 Brookings Dr., St. Louis, MO 63130, USA. E-mail: benschahary@wustl.edu

**Members of the natural resistance-associated macrophage protein (NRAMP) family are evolutionarily conserved metal ion transporters that play an essential role in regulating intracellular divalent cation homeostasis in both prokaryotes and eukaryotes. *Malvolio* (*Mvl*), the sole NRAMP family member in insects, plays a role in food choice behaviors in *Drosophila* and other species. However, the specific physiological and cellular processes that require the action of *Mvl* for appropriate feeding decisions remain elusive. Here, we show that normal food choice requires *Mvl* function specifically in the dopaminergic system, and can be rescued by supplementing food with manganese. Collectively, our data indicate that the action of the *Mvl* transporter affects food choice behavior via the regulation of dopaminergic innervation of the mushroom bodies, a principle brain region associated with decision-making in insects. Our studies suggest that the homeostatic regulation of the intraneuronal levels of divalent cations plays an important role in the development and function of the dopaminergic system and associated behaviors.**

Keywords: *D. melanogaster*, dopamine, food choice, fruit fly, ionomics, manganese, *Mvl*, NRAMP, Slc11a, vinegar fly

Received 21 October 2016, revised 16 February 2017, accepted for publication 16 February 2017

Correct food choices are essential for the survival and fitness of all animals. However, the genetic, neuronal and environmental factors that regulate these basal behaviors

are still not well understood. Previous studies in *Drosophila melanogaster* indicated that flies carrying a mutation in the gene *Malvolio* (*Mvl*), the insect homolog of the evolutionary-conserved protein family of natural resistance-associated macrophage proteins (NRAMPs) divalent metal ions transporters (D'Souza *et al.* 1999), lead to a reduced appetitive discrimination between high and low sugar-containing foods (Rodrigues *et al.* 1995). The effects of the *Mvl* mutation on food choice behavior could be rescued by supplementing food with manganese or iron (Orgad *et al.* 1998). In addition, studies in the honeybee *Apis mellifera* showed that expression levels of *Mvl* in the brain differ between bees, which perform different tasks, and are associated with changes in the behavioral response threshold to sugar stimuli in bees. Similarly, feeding manganese to young, pre-foraging bees leads to a lower response threshold to sugar stimuli, and a precocious transition to foraging behavior (Ben-Shahar *et al.* 2004; Søvik *et al.* 2015). The effect of the *Mvl* mutation on taste behavior is not because of abnormal gustatory sensory signaling. These data suggest that the effects of *Mvl* mutations on food choice are mediated via central elements of the feeding decision neuronal circuit or non-neuronal tissues (Rodrigues *et al.* 1995). The NRAMP genes are also expressed in the nervous systems of vertebrates, but their physiological roles in these tissues are not well understood (Evans *et al.* 2001; Ke *et al.* 2005; Skjorringe *et al.* 2015).

In addition to the genetic support for the role of *Mvl* in regulating feeding, other studies indicated that trace metals such as manganese and iron contribute to the regulation of organismal energy homeostasis via diverse mechanisms. For example, manganese can act as an insulin-mimetic *in vitro* and *in vivo* in mammalian models via unknown mechanisms (Nakai *et al.* 2005). Similarly, environmental exposure to manganese, iron and other trace metals has been implicated in the development of both *type I* and *type II* diabetes (Meyer & Spence 2009), and genetic variations in the human SLC11A1 (NRAMP1) gene are associated with the development of *type I* diabetes in humans (Yang *et al.* 2011). Independently, iron deficiency has been implicated in obesity (García *et al.* 2009; McClung & Karl 2009). Manganese and iron are also essential for mitochondrial respiration and cellular metabolism (Chen *et al.* 2015b; Levi & Rovida 2009; Mena *et al.* 2015; Mühlhoff *et al.* 2015; Pierrel *et al.* 2007). Together, these studies by others and us indicate that metal transporters play an important role in regulating diverse cognitive and metabolic functions in animals, and suggest that disruptions in the function of such transporters could lead to abnormal behavioral outcomes (Ávila *et al.* 2014; Dusek *et al.*

2015; Roels *et al.* 2012; Søvik *et al.* 2015; Su *et al.* 2015). Consequently, here, we used the power of *Drosophila* genetics to investigate the cellular and molecular pathways that may explain the effects of *Mvl* on behavioral feeding decisions and organismal metabolic homeostasis.

## Materials and methods

### Fly stocks and maintenance

Flies were reared on standard *Drosophila* cornmeal medium at 25°C and 60% humidity under a 12h:12h light/dark cycle. GAL4 lines, such as UAS-*Dcr2*, UAS-*Mvl*-RNAi and UAS-tetanus toxin (TNT) lines, were from the Bloomington *Drosophila* Stock Center (Bloomington, IN, USA).

### Behavioral experiments

The single-dye feeding decisions paradigm was as previously described by us and others (Lu *et al.* 2012; Orgad *et al.* 1998). In short, the 'feeding error index' represents the proportion of flies with red abdomens relative to total number of flies in a single assay. As single-dye feeding assays cannot discriminate between flies that made a correct feeding choice and flies that did not eat at all (both will have no dye in the abdomen), we also measured the effects of genotypes and treatments on the general feeding drive of flies by allowing groups of animals to feed on plates in which all wells had 10 mM trehalose with the same red dye as in feeding choice assays.

### Determination of the elemental composition of fly tissues by ICP-MS

Four independent samples of 10 pooled fly heads per genotype were analyzed for B, Na, Mg, Al, P, S, K, Ca, Mn, Fe, Co, Cu, Zn, As, Se, Rb, Mg and Cd. Inductively coupled plasma mass spectrometry (ICP-MS) analyses followed the standard Baxter laboratory protocol as previously described (Ziegler *et al.* 2013).

### Quantification of biogenic amines from fly brains using high-pressure liquid chromatography

Adult fly brains were dissected in chilled PBS. Samples consisting of 10 pooled brains were flash-frozen in liquid nitrogen and stored at  $-80^{\circ}\text{C}$  until analysis. Biogenic amine content was extracted in 15  $\mu\text{l}$  of 0.2 M perchloric acid containing 10  $\mu\text{g}/\mu\text{l}$  of the high-pressure liquid chromatography (HPLC) standard dihydroxybenzylamine (Sigma-Aldrich, St. Louis, MO, USA) as previously described (Scheiner *et al.* 2014; Søvik *et al.* 2013). The supernatant of each sample separated across an HR-80 column (ESA Biosciences, Inc., Chelmsford, MA, USA) using an Agilent 1200 Series HPLC system (Agilent Technologies, Santa Clara, CA, USA). Levels of biogenic amines were quantified using an ESA Coulochem III electrochemical detector with an ESA 5011A analytical cell. Standard curves for octopamine, dopamine and serotonin (Sigma-Aldrich) were used to calculate the total amount of each amine in the tested samples.

### Quantification of mRNA expression levels by real-time qRT-PCR

Total RNA was extracted from pools of approximately 30 adult heads using the TRIzol reagent (Thermo Fisher Scientific, Waltham, MA USA). First strand cDNA was prepared using *SuperScript* II reverse transcriptase (Thermo Fisher Scientific, Waltham, MA USA). Gene-specific assays were conducted with a SYBR Green kit on a 7500 real-time PCR System (Thermo Fisher Scientific, Waltham, MA USA). Each cDNA sample was run in triplicate PCR reactions. The housekeeping gene *rp49* was used as a loading control. Relative expression data were analyzed and presented as expression

fold-difference by using the  $\Delta\Delta\text{CT}$  method, as we have previously described (Lu *et al.* 2012; Lu *et al.* 2014; Søvik *et al.* 2015; Zheng *et al.* 2014).

### Immunohistochemistry

Antibody staining of dopaminergic neurons was as previously described (Lu *et al.* 2012; Lu *et al.* 2014; Zelle *et al.* 2013; Zheng *et al.* 2014). Brains were co-stained with anti-GFP (Thermo Fisher Scientific, Waltham, MA USA) and nc82 antibodies (Developmental Studies Hybridoma Bank, Iowa City, IA, USA), and mounted using the VECTASHIELD HardSet Antifade Mounting Medium with DAPI (Vector Laboratories, Burlingame, CA, USA). Imaging was performed using Nikon A1 confocal microscope. Figures were generated from maximal intensity Z-stacks using the Nikon NIS-Elements software package.

### Statistical analyses

Feeding behavioral data and brain ionomics were analyzed with Student's *t*-tests (StatPlus, AnalystSoft, WALNUT, CA USA). All values for the statistical tests are described in Tables S1–S5, Supporting Information. Biogenic amines levels were analyzed by a linear mixed model with genotype as a categorical factor. The HPLC analysis batch was controlled for in the model as well. Dry weight and glucose levels were analyzed by one-way analysis of variance followed by a Tukey *post hoc* test ( $P < 0.05$ ).


## Results

### *Mvl* is required in neuronal tissues for normal food choice behavior

Analysis of the spatial *Mvl* mRNA expression data from the FlyAtlas database (Robinson *et al.* 2013) indicated that it is broadly expressed throughout the adult body (Fig. 1a). Therefore, to identify the specific tissues in which *Mvl* is required for feeding decisions, we employed a tissue-specific RNAi-based gene knockdown approach (Brand & Perrimon 1993). To test whether the available UAS-*Mvl*-RNAi line can effectively knockdown endogenous mRNAs, we first showed the effect of the UAS-*Mvl*-RNAi on the global expression of *Mvl* with the ubiquitous  $\alpha\text{Tub84B}$ -GAL4 line (Fig. 1b).

Previous studies indicated that the abnormal feeding phenotype in *Mvl* mutant flies can be rescued on a physiological timescale with manganese or iron food supplements (Orgad *et al.* 1998). However, because *Mvl* is broadly expressed throughout development (not shown), it is still impossible to know whether the behavioral phenotype of adult mutants is due, at least in part, to abnormal developmental processes. Therefore, we investigated the impact of global *Mvl* knockdown on feeding decisions specifically in the adult stage by expressing the UAS-*Mvl*-RNAi with the conditional heat-induced *hsp70*-GAL4 driver. We found that under heat-shock conditions, adult flies exhibited abnormal feeding behavior (Fig. 1c and Tables S1 and S2). These data further suggest that the effects of the *Mvl* gene knockdown on behavior are not solely dependent on post-embryonic development.

Because *Mvl* is broadly expressed (Fig. 1a), we next used specific organ-enriched GAL4 drivers to investigate whether the effects of *Mvl* knockdown on food choice behavior are tissue-specific. The tissue-level RNAi screen showed that




**Figure 1: Tissue-specific role for *Mvl* in feeding decisions.** (a) Tissue distribution of *Mvl* expression in adult *Drosophila*. Data are from FlyAtlas (Robinson *et al.* 2013). Error bars represent SEM ( $n=4$  independent microarrays). (b) Global ectopic expression of transgenic *Mvl*-RNAi with the *tubulin* GAL4 driver reduces *Mvl* expression. (c) Conditional *Mvl* gene knockdown in the adult stage with the heat-shock-induced *Hsp70* GAL4 driver ('-', without heat shock; '+', with heat shock). One-way ANOVA; \* $P < 0.05$ . (d) General feeding-drive control experiment. Genotypes and conditions as in (c). (e) Tissue-specific *Mvl* knockdown. (f) General feeding-drive control experiment. Genotypes and conditions as in e (see Table S1 for statistics).

*Mvl* expression in the nervous system (*elav*-GAL4) is necessary for normal food choice behavior (Fig. 1d). In contrast, expressing the *Mvl*-RNAi with GAL4 drivers enriched in the fat body (*ppl*), hemocytes and fat body (*cg*), salivary glands and fat body (*R4*), gut (*drm*), glia (*repo*) or everywhere outside the nervous system (*DJ761*), did not affect feeding decisions (Fig. 1d and Table S1). Together, these data indicate that normal feeding decisions require *Mvl* activity in the nervous system. We did not observe any effects of the *Mvl* gene knockdown on general feeding drive in any of the tested genotypes (Fig. 1f and Table S2).

**Effect of neuronal *Mvl* knockdown on behavior can be rescued by manganese**

Previous work showed that manganese supplements can rescue the effect of the *Mvl* mutation on feeding decisions (Orgad *et al.* 1998). Therefore, we next asked whether the effect of neuronal-specific *Mvl* knockdown on feeding behavior could be rescued by a manganese supplement as well. We found that supplementing fly food with 5 mM  $Mn^{2+}$  was sufficient to rescue the effects of the neuronal *Mvl*-RNAi downregulation on feeding decisions (Fig. 2a). However,



**Figure 2: Manganese food supplement rescues effects of neuronal *Mvl* knockdown on behavior.** (a) Feeding manganese rescues the effect of neuronal *Mvl* knock down on food choice behavior. \*\*, *t*-test,  $t_{15} = 5.1628$ ,  $P < 0.01$ . (b) General feeding-drive control experiment. Genotypes and conditions as in (a). (c) Effect of neuronal *Mvl* knockdown and manganese treatment on total dry weight. Different letters above bars represent significantly different groups (ANOVA,  $P < 0.05$ , Tukey *post hoc* test). (d) Effect of neuronal *Mvl* knockdown and manganese on glucose levels (ANOVA, NS).

neither genetic background nor  $Mn^{2+}$  treatment had an effect on the overall feeding drive of adult flies (Fig. 2b). We also found that the  $Mn^{2+}$  supplement rescued a small, but significant, decrease in total dry body weight observed in flies with the pan-neuronal *Mvl* knockdown (Fig. 2c), but not in total glucose content (Fig. 2d), which possibly reflects the increased consumption of non-nutritive food in *Mvl* knock-down flies. Together, our data suggest that *Mvl*-dependent regulation of manganese in neuronal tissues plays a role in food choice behaviors in the fly. These data also indicate that manganese, and possibly other divalent cations, can enter cells via alternative, NRAMP-independent pathways as has been reported in other model systems (Chen *et al.* 2015a; Garcia-Rodriguez *et al.* 2015; Jenkins *et al.* 2016).


### Effects of the neuronal *Mvl* knockdown on the head ionome

Because *Mvl* is a divalent metal ion transporter, we next examined the elemental composition of fly heads in *elav>Mvl-RNAi* flies relative to wild-type controls. Surprisingly, we found that levels of  $Mn^{2+}$ ,  $Fe^{2+}$  and  $Cu^{2+}$ , which are ions known to be transported by *Mvl* (Ben-Shahar

*et al.* 2004; Betti *et al.* 2011; Folwell *et al.* 2006; Orgad *et al.* 1998; Southon *et al.* 2008), were not affected by the neuronal *Mvl* knockdown (Fig. 2 and Table S3). In contrast, levels of  $Na^+$ ,  $K^+$  and  $Rb^+$ , three elements known to have similar chemical properties, showed a small but significant reduction in the *Mvl* knockdown animals (Fig. 3 and Table S3). These data indicate that the neuronal-specific knockdown of *Mvl* is not sufficient to impact the overall metal homeostasis of trace metals, possibly because levels of these metals in the hemolymph and non-neuronal tissues in the head mask neuronal-specific changes. Whether the small effects of the neuronal *Mvl* knockdown on the levels of other ions are directly related to the transporter activity of *Mvl*, or whether they represent indirect, downstream effects of the knockdown on other ionic transport pathways was not investigated further in this study.

### *Mvl* expression in the dopaminergic system is required for normal food choice behavior

Feeding decisions in all animals, including flies, are driven by complex neuronal circuits, and depend on the integration of diverse internal and external signals (Albin *et al.* 2015; Flood *et al.* 2013; Huetteroth *et al.* 2015; Itskov & Ribeiro 2013; LeDue *et al.* 2015; Perisse *et al.* 2016; Yapici *et al.* 2016). To further narrow the possible neuronal populations that require *Mvl* function for normal feeding decisions, we next drove the *Mvl-RNAi* construct with GAL4 drivers that are active in specific neuronal subpopulation that have been previously reported to play a role in feeding decisions. Our screen included drivers specific to gustatory receptor (*Poxn*) neurons (Dahanukar *et al.* 2001), dopaminergic (*TH*) neurons (Marella *et al.* 2012; Melcher & Pankratz 2005), octopaminergic (*Tdc2*) neurons (Zhang *et al.* 2013), cholinergic (*Cha*) neurons (Yapici *et al.* 2016), neuropeptidergic (*dimm*) neurons (Kahsai *et al.* 2010; Taghert & Nitabach 2012; Zhang *et al.* 2013), *Neuropeptide F*-expressing (*Npf*) neurons (Shen & Cai 2001), *Adipokinetic hormone*-expressing (*Akh*) neurons (Kim & Neufeld 2015) and the circadian system (*Tim*) (DiAngelo *et al.* 2011). This limited screen showed that *Mvl* knockdown specifically in dopaminergic neurons (*TH-GAL4*; Fig. 4a and Table S4) phenocopied the abnormal food choice phenotype we observed in animals with pan-neuronal *Mvl* knockdown (Fig. 1e). In contrast, expression of the *Mvl-RNAi* in all gustatory receptor neurons (*Poxn*) or neurons that express the neuropeptide *adipokinetic hormone* (*Akh*), which have been previously implicated in the regulation of feeding and metabolism (Itskov & Ribeiro 2013; Pool & Scott 2014), showed a small but significant decrease in feeding error rates relative to wild-type controls (Fig. 4a and Table S4). Unexpectedly, these data indicate that *Mvl* knockdown has different effects on feeding decisions in different neuronal elements of the feeding decision circuit. Regardless, *Mvl* gene knockdown in any of the neuronal populations we have examined did not affect the general feeding drive of adult flies, further confirming that the effects observed represents an abnormal decision-making phenotype rather than simply observing the behavior of unhealthy flies (Fig. 4b and Table S5).



**Figure 3: Effect of neuronal *Mvl* knockdown on the head ionome.** The elemental metal composition of fly heads in neuronal *Mvl* knockdown and wild-type control flies ( $n=4$ ; \*,  $P < 0.05$ ; see Table S3 for statistics).


### Effects of neuronal *Mvl* knockdown on behavior are not mediated via overall changes in biogenic amine synthesis

Because several of the metal ions that were previously shown to be transported by *Mvl* function as cofactors in neurotransmitter synthesis, and because of the effect of *Mvl* knockdown in dopaminergic neurons on food choice behavior, we next investigated the effects of *Mvl*-RNAi expression on the syntheses of biogenic amines in the fly brain. Using HPLC, we measured levels of dopamine, serotonin and octopamine from the brains of pan-neuronal *Mvl* knockdown and wild-type flies. Our data show that the overall brain levels of the three primary biogenic amines in the insect brain, dopamine, octopamine and serotonin (Evans 1980; Mesce 2002), were not affected by neuronal *Mvl* knockdown (Fig. 5a). These data suggest that the effects of *Mvl* knockdown on behavior are not a direct consequence of dramatic physiological changes in the brain biogenic amines systems.

Since *Mvl* knockdown in dopaminergic neurons does not seem to affect overall dopamine synthesis, we next asked whether the effects of dopaminergic *Mvl* knockdown are because of a reduced dopamine release. Previous studies indicated that disruptions of the dopaminergic signaling pathway can lead to a general low appetite (hypophagia) (Krashes *et al.* 2009; Riemensperger *et al.* 2011). In agreement with the earlier findings, we found that blocking dopamine release by the ectopic expression of the TNT with the dopaminergic *TH*-GAL4 driver resulted in general hypophagia (Fig. 5b,c), which is unlike the phenotype we observed in *Mvl* knockdown animals. We also found that the mRNA levels of *ple* (the gene encoding *TH*) were not affected by the expression of the *Mvl*-RNAi in the dopaminergic system. Together, these

data further indicate that the effects of *Mvl* knockdown in the dopaminergic system on food choice behavior are not likely to be directly mediated via modulation of dopamine synthesis or release (Fig. 5d).

Since *Mvl* activity does not seem to directly affect dopamine synthesis or release, we also examined the effects of the *Mvl* knockdown on dopaminergic neuron morphology. As in other animals, the dopaminergic system in the fly is complex and heterogeneous. Previous studies have identified the protocerebral anterior medial neuronal cluster, which innervates the mushroom bodies, as the primary modulatory pathways for the integration of food-rewards in long-term memory (Huetteroth *et al.* 2015; Liu *et al.* 2012). Independently, the activity of TH-VUM, a single dopaminergic neuron that innervates the subesophageal ganglion (SOG) taste center, was shown to regulate the appetitive acceptance of sugar rewards (Marella *et al.* 2012). To test whether *Mvl* knockdown affected dopaminergic morphology, we expressed a membrane tethered GFP (UAS-CD8::GFP) with the *TH*-GAL4 line, with or without the UAS-*Mvl*-RNAi. These studies showed that *Mvl* knockdown animals exhibit a dramatic reduction in the dopaminergic innervation of both the mushroom bodies and the SOG (Figs. 5f and S2b). By contrast, control animals show extensive innervation of the mushroom bodies in the protocerebrum and TH-VUM signal in the SOG (Figs. 5e and S2a). To further establish a causal link between the action of *Mvl* in the dopaminergic system and food-choice behaviors, we also attempted to study dopaminergic neuronal morphology and food-choice behavior in flies carrying the previously published *Mvl*<sup>97f</sup> allele (D'Souza *et al.* 1999; Orgad *et al.* 1998; Rodrigues *et al.* 1995). However, in our hands, the observed phenotype of flies carrying this allele in two different genetic backgrounds



**Figure 4: Effect of *Mvl* knock-down in specific neuronal populations.** (a) Feeding choice assay. GAL4 drivers: *Poxn*, all gustatory receptor neurons; *TH*, dopaminergic neurons; *Tdc2*, octopaminergic neurons; *Cha*, cholinergic neurons; *dimm*, neuropeptidergic neurons; *Npf*, Neuropeptide F neurons; *Akh*, Adipokinetic hormone neurons; *Tim*, circadian neurons. \*\*, p < 0.01. (b) General feeding-drive control experiment. Genotypes as in (a). See Tables S4 and S5 for statistics.

was weak and inconsistent (not shown). Nevertheless, our findings suggest that *Mvl* activity in dopaminergic neurons play a critical role in modulating synaptic connectivity in the dopaminergic system, which possibly affects specific neuronal pathways associated with food-choice decisions.


## Discussion

In recent years, studies of the dopaminergic system across animal phyla have indicated that it plays a conserved, important role in regulating cognitive processes associated with decision-making in general, and feeding decisions in particular (Barron *et al.* 2015; de Bivort & van Swinderen 2016; Khani & Rainer 2016; Stopper & Floresco 2015). Consequently, the highly conserved cellular and biochemical properties of the dopaminergic system across mammals and invertebrates indicate that mechanistic insights gained here are likely relevant to neuronal and behavioral functions of metal transporters in many other animal species, including humans.

We have previously showed that chronic feeding of wild-type flies and honey bees with manganese increased the overall brain levels of dopamine, and adversely affected bee foraging decisions (Søvik *et al.* 2015). We have interpreted these published data to suggest that the effects

of environmental exposure to manganese were mediated via a direct effect on the dopaminergic synthesis pathway. Therefore, we originally expected that the effects of knocking down *Mvl*, an established manganese transporter, on feeding behavior will be associated with changes in dopaminergic levels as well. Our finding that the adverse behavioral outcomes of dopaminergic *Mvl* knockdown are not associated with overall changes in dopamine levels could be possibly explained by the insensitivity of currently available assays to very small changes in dopamine levels. Alternatively, these data could indicate that the interaction between the dopaminergic system and metal ion homeostasis is more complex than we originally expected, and may independently affect brain functions at both the developmental and physiological timescales. Nonetheless, our finding that *Mvl* knockdown in dopaminergic neurons changes their innervation patterns further supports this alternative interpretation.

Previous studies showed that the dopaminergic system in the fly brain is remodeled at the pupal stage during metamorphosis (Budnik & White 1988). However, we still do not know whether the effect of *Mvl* knockdown on the morphology and innervation patterns of dopaminergic neurons is because of its requirement at the pupal neuronal remodeling stage, or whether it reflects an interaction between *Mvl* activity and post-pupal neuronal plasticity. In addition, although the focus of this study



**Figure 5: Effects of *Mvl* knock-down on the dopaminergic system.** (a) HPLC analyses of biogenic amines levels. Dopamine,  $t_{34} = -0.9463$ , NS; Octopamine,  $t_{34} = -0.8967$ , NS; Serotonin,  $t_{31} = -0.4124$ , NS. (b) Effect of blocking dopaminergic synaptic release with the ectopic expression of the tetanus toxin on feeding errors ( $t_{21} = 4.8833$ , \*,  $P < 0.001$ ). (c) Effect of blocking dopaminergic synaptic release with the ectopic expression of the tetanus toxin on feeding rate ( $t_{11} = 2.668$ , \*,  $P = 0.02187$ ). (d) Effect of dopaminergic *Mvl* knock-down on mRNA expression levels of *ple* (*TH>Mvl-RNAi*). (e) Dopaminergic projection patterns in wild-type brains (*TH-GAL4>UAS-CD8::GFP*). (f) Effect of dopaminergic *Mvl* knockdown on dopaminergic neuronal projection patterns (*TH-GAL4>UAS-CD8::GFP//UAS-Mvl-RNAi*). In (e) and (f), solid arrowheads mark innervations of the mushroom bodies, and empty arrowhead mark innervations of the subesophageal ganglion.

is on the interaction between *Mvl* and the dopaminergic system, we cannot exclude a possible effect of *Mvl* knockdown on the morphology of other neuronal classes, and therefore, on other, dopamine-independent behavioral phenotypes.

We found that knocking down *Mvl* expression specifically in the dopaminergic system is sufficient to phenocopy the pan-neuronal knockdown phenotype. We interpret these data to suggest that, in agreement with previous studies, the dopaminergic system acts as a 'master switch', which determines the probability that an animal will respond behaviorally to an appetitive stimulus. The concept of a dopaminergic 'master switch' is further supported by the contrary effects of *Mvl* knockdown specifically in the gustatory sensory system and *Npf*-expressing neurons, which resulted in an increase in the appetitive preference for sugar relative to control animals. However, to completely

understand how *Mvl*-dependent changes in dopaminergic innervation affect feeding-related decision-making behaviors will require the identification and characterization of the relevant neuronal elements that receive inputs from *Mvl*-expressing dopaminergic neurons. Nevertheless, these data suggest that *Mvl*, and its possible effects on metal ion homeostasis, play independent roles in different elements of the neuronal circuit associated with feeding decisions.

Data presented here strongly support the idea that metal ion homeostasis in specific neuronal circuits could have a dramatic impact on neuronal functions and behavior. The conserved biochemical properties of the molecules involved in metal ion homeostasis across distant phylogenetic clades indicate that insights gained in the current studies are likely conserved in other animal species, including humans.

## References

- Albin, S.D., Kaun, K.R., Knapp, J.-M., Chung, P., Heberlein, U. & Simpson, J.H. (2015) A subset of serotonergic neurons evokes hunger in adult *Drosophila*. *Curr Biol* **25**, 2435–2440.
- Ávila, D., Puntel, R., Folmer, V., Rocha, J., dos Santos, A. & Aschner, M. (2014) Manganese neurotoxicity. In Kostrzewa, R.M. (ed), *Handbook of Neurotoxicity*. Springer, New York, pp. 843–864.
- Barron, A.B., Gurney, K.N., Meah, L.F., Vasilaki, E. & Marshall, J.A. (2015) Decision-making and action selection in insects: inspiration from vertebrate-based theories. *Front Behav Neurosci* **9**, 216.
- Ben-Shahar, Y., Dudek, N.L. & Robinson, G.E. (2004) Phenotypic deconstruction reveals involvement of manganese transporter malvolio in honey bee division of labor. *J Exp Biol* **207**, 3281–3288.
- Bettedi, L., Aslam, M.F., Szular, J., Mandilaras, K. & Missirlis, F. (2011) Iron depletion in the intestines of Malvolio mutant flies does not occur in the absence of a multicopper oxidase. *J Exp Biol* **214**, 971–978.
- de Bivort, B.L. & van Swinderen, B. (2016) Evidence for selective attention in the insect brain. *Curr Opin Insect Sci* **15**, 9–15.
- Brand, A.H. & Perrimon, N. (1993) Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* **118**, 401–415.
- Budnik, V. & White, K. (1988) Catecholamine-containing neurons in *Drosophila melanogaster*: distribution and development. *J Comp Neurol* **268**, 400–413.
- Chen, P., Chakraborty, S., Mukhopadhyay, S., Lee, E., Paoliello, M.M., Bowman, A.B. & Aschner, M. (2015a) Manganese homeostasis in the nervous system. *J Neurochem* **134**, 601–610.
- Chen, P., Chakraborty, S., Peres, T.V., Bowman, A.B. & Aschner, M. (2015b) Manganese-induced neurotoxicity: from *C. elegans* to humans. *Toxicol Res* **4**, 191–202.
- Dahanukar, A., Foster, K., van der Goes van Naters, W.M. & Carlson, J.R. (2001) A Gr receptor is required for response to the sugar trehalose in taste neurons of *Drosophila*. *Nat Neurosci* **4**, 1182–1186.
- DiAngelo, J.R., Erion, R., Crocker, A. & Sehgal, A. (2011) The central clock neurons regulate lipid storage in *Drosophila*. *PLoS One* **6**, e19921.
- D'Souza, J., Cheah, P.Y., Gros, P., Chia, W. & Rodrigues, V. (1999) Functional complementation of the malvolio mutation in the taste pathway of *Drosophila melanogaster* by the human natural resistance-associated macrophage protein 1 (Nramp-1). *J Exp Biol* **202**, 1909–1915.
- Dusek, P., Roos, P.M., Litwin, T., Schneider, S.A., Flaten, T.P. & Aaseth, J. (2015) The neurotoxicity of iron, copper and manganese in Parkinson's and Wilson's diseases. *J Trace Elem Med Biol* **31**, 193–203.
- Evans, P.D. (1980) Biogenic amines in the insect nervous system. In Berridge, M.J., Treherne, J.E. & Wigglesworth, V.B. (eds), *Advances in Insect Physiology*. Academic Press, pp. 317–473.
- Evans, C., Harbuz, M. & Ostenfeld, T. (2001) Nramp1 is expressed in neurones and is associated with behavioural and immune responses to stress. *Neurogenetics* **3**, 69–78.
- Flood, T.F., Iguchi, S., Gorczyca, M., White, B., Ito, K. & Yoshihara, M. (2013) A single pair of interneurons commands the *Drosophila* feeding motor program. *Nature* **499**, 83–87.
- Folwell, J.L., Barton, C.H. & Shepherd, D. (2006) Immunolocalisation of the *D. melanogaster* Nramp homologue Malvolio to gut and Malpighian tubules provides evidence that Malvolio and Nramp2 are orthologous. *J Exp Biol* **209**, 1988–1995.
- García, O.P., Long, K.Z. & Rosado, J.L. (2009) Impact of micronutrient deficiencies on obesity. *Nutr Rev* **67**, 559–572.
- García-Rodríguez, N., Manzano-Lopez, J., Muñoz-Bravo, M., Fernández-García, E., Muniz, M. & Wellinger, R.E. (2015) Manganese redistribution by calcium-stimulated vesicle trafficking bypasses the need for P-type ATPase function. *J Biol Chem* **290**, 9335–9347.
- Huetteroth, W., Perisse, E., Lin, S., Klappenbach, M., Burke, C. & Waddell, S. (2015) Sweet taste and nutrient value subdivide rewarding dopaminergic neurons in *Drosophila*. *Curr Biol* **25**, 751–758.
- Itskov, P.M. & Ribeiro, C. (2013) The dilemmas of the gourmet fly: the molecular and neuronal mechanisms of feeding and nutrient decision making in *Drosophila*. *Front Neurosci* **7**, 12.
- Jenkins, J., Papkovsky, D.B. & Dmitriev, R.I. (2016) The Ca<sup>2+</sup>/Mn<sup>2+</sup>-transporting SPCA2 pump is regulated by oxygen and cell density in colon cancer cells. *Biochem J* **473**, 2507–2518.
- Kahsai, L., Kapan, N., Dircksen, H., Winther, A.M. & Nassel, D.R. (2010) Metabolic stress responses in *Drosophila* are modulated by brain neurosecretory cells that produce multiple neuropeptides. *PLoS One* **5**, e11480.
- Ke, Y., Chang, Y.Z., Duan, X.L., Du, J.R., Zhu, L., Wang, K., Yang, X.D., Ho, K.P. & Qian, Z.M. (2005) Age-dependent and iron-independent expression of two mRNA isoforms of divalent metal transporter 1 in rat brain. *Neurobiol Aging* **26**, 739–748.
- Khani, A. & Rainer, G. (2016) Neural and neurochemical basis of reinforcement-guided decision making. *J Neurophysiol* **116**, 724–741.
- Kim, J. & Neufeld, T.P. (2015) Dietary sugar promotes systemic TOR activation in *Drosophila* through AKH-dependent selective secretion of Dilp3. *Nat Commun* **6**, 6846.
- Krashes, M.J., DasGupta, S., Vreede, A., White, B., Armstrong, J.D. & Waddell, S. (2009) A neural circuit mechanism integrating motivational state with memory expression in *Drosophila*. *Cell* **139**, 416–427.
- LeDue, E.E., Chen, Y.C., Jung, A.Y., Dahanukar, A. & Gordon, M.D. (2015) Pharyngeal sense organs drive robust sugar consumption in *Drosophila*. *Nat Commun* **6**, 6667.
- Levi, S. & Rovida, E. (2009) The role of iron in mitochondrial function. *Biochim Biophys Acta* **1790**, 629–636.
- Liu, C., Placais, P.-Y., Yamagata, N., Pfeiffer, B.D., Aso, Y., Friedrich, A.B., Siwanowicz, I., Rubin, G.M., Preat, T. & Tanimoto, H. (2012) A subset of dopamine neurons signals reward for odour memory in *Drosophila*. *Nature* **488**, 512–516.
- Lu, B., LaMora, A., Sun, Y., Welsh, M.J. & Ben-Shahar, Y. (2012) ppk23-Dependent chemosensory functions contribute to courtship behavior in *Drosophila melanogaster*. *PLoS Genet* **8**, e1002587.
- Lu, B., Zelle, K.M., Seltzer, R., Hefetz, A. & Ben-Shahar, Y. (2014) Feminization of pheromone-sensing neurons affects mating decisions in *Drosophila* males. *Biology Open* **3**, 152–160.
- Marella, S., Mann, K. & Scott, K. (2012) Dopaminergic modulation of sucrose acceptance behavior in *Drosophila*. *Neuron* **73**, 941–950.
- McClung, J.P. & Karl, J.P. (2009) Iron deficiency and obesity: the contribution of inflammation and diminished iron absorption. *Nutr Rev* **67**, 100–104.
- Melcher, C. & Pankratz, M.J. (2005) Candidate gustatory interneurons modulating feeding behavior in the *Drosophila* brain. *PLoS Biol* **3**, e305.
- Mena, N.P., Urrutia, P.J., Lourido, F., Carrasco, C.M. & Núñez, M.T. (2015) Mitochondrial iron homeostasis and its dysfunctions in neurodegenerative disorders. *Mitochondrion* **21**, 92–105.
- Mesce, K.A. (2002) Metamodulation of the biogenic amines: second-order modulation by steroid hormones and amine cocktails. *Brain Behav Evol* **60**, 339–349.
- Meyer, J.A. & Spence, D.M. (2009) A perspective on the role of metals in diabetes: past findings and possible future directions. *Metallomics* **1**, 32.
- Mühlenhoff, U., Hoffmann, B., Richter, N., Rietzschel, N., Spantgar, F., Stehling, O., Uzarska, M.A. & Lill, R. (2015) Compartmentalization of iron between mitochondria and the cytosol and its regulation. *Eur J Cell Biol* **94**, 292–308.
- Nakai, M., Sekiguchi, F., Obata, M., Ohtsuki, C., Adachi, Y., Sakurai, H., Orvig, C., Rehder, D. & Yano, S. (2005) Synthesis and insulin-mimetic activities of metal complexes with 3-hydroxypyridine-2-carboxylic acid. *J Inorg Biochem* **99**, 1275–1282.



- Orgad, S., Nelson, H., Segal, D. & Nelson, N. (1998) Metal ions suppress the abnormal taste behavior of the *Drosophila* mutant malvolio. *J Exp Biol* **201**, 115–120.
- Perisse, E., Oswald, D., Barnstedt, O., Talbot, C.B., Huetteroth, W. & Waddell, S. (2016) Aversive learning and appetitive motivation toggle feed-forward inhibition in the *Drosophila* mushroom body. *Neuron* **90**, 1086–1099.
- Pierrel, F., Cobine, P.A. & Winge, D.R. (2007) Metal ion availability in mitochondria. *Biometals* **20**, 675–682.
- Pool, A.H. & Scott, K. (2014) Feeding regulation in *Drosophila*. *Curr Opin Neurobiol* **29**, 57–63.
- Riemensperger, T., Isabel, G., Coulom, H., Neuser, K., Seugnet, L., Kume, K., Iche-Torres, M., Cassar, M., Strauss, R., Preat, T., Hirsh, J. & Birman, S. (2011) Behavioral consequences of dopamine deficiency in the *Drosophila* central nervous system. *Proc Natl Acad Sci USA* **108**, 834–839.
- Robinson, S.W., Herzyk, P., Dow, J.A. & Leader, D.P. (2013) FlyAtlas: database of gene expression in the tissues of *Drosophila melanogaster*. *Nucleic Acids Res* **41**, D744–D750.
- Rodrigues, V., Cheah, P.Y., Ray, K. & Chia, W. (1995) malvolio, the *Drosophila* homologue of mouse NRAMP-1 (Bcg), is expressed in macrophages and in the nervous system and is required for normal taste behaviour. *EMBO J* **14**, 3007–3020.
- Roels, H.A., Bowler, R.M., Kim, Y., Claus Henn, B., Mergler, D., Hoet, P., Gocheva, V.V., Bellinger, D.C., Wright, R.O., Harris, M.G., Chang, Y., Bouchard, M.F., Riojas-Rodriguez, H., Menezes-Filho, J.A. & Tellez-Rojo, M.M. (2012) Manganese exposure and cognitive deficits: a growing concern for manganese neurotoxicity. *Neurotoxicology* **33**, 872–880.
- Scheiner, R., Toteva, A., Reim, T., Søvik, E. & Barron, A.B. (2014) Differences in the phototaxis of pollen and nectar foraging honey bees are related to their octopamine brain titers. *Front Physiol* **5**, 1–8.
- Shen, P. & Cai, H.N. (2001) *Drosophila* neuropeptide F mediates integration of chemosensory stimulation and conditioning of the nervous system by food. *J Neurobiol* **47**, 16–25.
- Skjorringe, T., Burkhart, A., Johnsen, K.B. & Moos, T. (2015) Divalent metal transporter 1 (DMT1) in the brain: implications for a role in iron transport at the blood-brain barrier, and neuronal and glial pathology. *Front Mol Neurosci* **8**, 19.
- Southon, A., Farlow, A., Norgate, M., Burke, R. & Camakaris, J. (2008) Malvolio is a copper transporter in *Drosophila melanogaster*. *J Exp Biol* **211**, 709–716.
- Søvik, E., Perry, C.J., LaMora, A., Barron, A.B. & Ben-Shahar, Y. (2015) Negative impact of manganese on honeybee foraging. *Biol Lett* **11**, 20140989.
- Søvik, E., Cornish, J.L. & Barron, A.B. (2013) Cocaine tolerance in honey bees. *PLoS One* **8**, e64920.
- Stopper, C.M. & Floresco, S.B. (2015) Dopaminergic circuitry and risk/reward decision making: implications for schizophrenia. *Schizophr Bull* **41**, 9–14.
- Su, C., Chen, K., Zou, Y., Shen, Y., Xia, B., Liang, G., Lv, Y., Wang, F., Huang, D. & Yang, X. (2015) Chronic exposure to manganese sulfate leads to adverse dose-dependent effects on the neurobehavioral ability of rats. *Environ Toxicol* **31**, 1571–1579.
- Taghert, P.H. & Nitabach, M.N. (2012) Peptide neuromodulation in invertebrate model systems. *Neuron* **76**, 82–97.
- Yang, J.H.M., Downes, K., Howson, J.M.M., Nutland, S., Stevens, H.E., Walker, N.M. & Todd, J.A. (2011) Evidence of association with type 1 diabetes in the SLC11A1 gene region. *BMC Med Genet* **12**, 59.
- Yapici, N., Cohn, R., Schusterreiter, C., Ruta, V. & Vosshall, L.B. (2016) A taste circuit that regulates ingestion by integrating food and hunger signals. *Cell* **165**, 715–729.
- Zelle, K.M., Lu, B., Pyfrom, S.C. & Ben-Shahar, Y. (2013) The genetic architecture of degenerin/epithelial sodium channels in *Drosophila*. *G3* **3**, 441–450.
- Zhang, T., Branch, A. & Shen, P. (2013) Octopamine-mediated circuit mechanism underlying controlled appetite for palatable food in *Drosophila*. *Proc Natl Acad Sci USA* **110**, 15431–15436.
- Zheng, X., Valakh, V., DiAntonio, A. & Ben-Shahar, Y. (2014) Natural antisense transcripts regulate the neuronal stress response and excitability. *Elife* **3**, e01849.
- Ziegler, G., Terauchi, A., Becker, A., Armstrong, P., Hudson, K. & Baxter, I. (2013) Ionomic screening of field-grown soybean identifies mutants with altered seed elemental composition. *Plant Genome* **6**. doi:10.3835/plantgenome2012.070012

## Acknowledgments

This study was funded by a Children's Discovery Institute grant MD-II-2009-170 to J.G.D. and Y.B.-S., NIH R21NS089834 to Y.B.-S. We thank members of the Ben-Shahar laboratory for their helpful comments on earlier versions of the manuscript. We thank Ivan Baxter and Greg Ziegler of the USDA-ARS Ionomic Profiling Facility at the Donald Danforth Plant Science Center for assistance with elemental analysis.

## Supporting Information

Additional supporting information may be found in the online version of this article at the publisher's web-site:

**Table S1:** Statistical analyses of % food choice error in *Mvl* knockdown vs. wild type flies (Fig. 1c–f).

**Table S2:** Statistical analyses of % feeding controls in *Mvl* knockdown vs. wt flies (Fig. 1c–f).

**Table S3:** Effect of neuronal *Mvl* knockdown on elemental composition of fly heads (Fig. 3).

**Table S4:** Statistical analyses of % food choice error in *Mvl* knockdown vs. wt flies (Fig. 4a).

**Table S5:** Statistical analyses of % feeding controls in *Mvl* knockdown vs. wt flies (Fig. 4b).